

# UC San Diego

## UC San Diego Previously Published Works

### Title

Correction: Broad dengue neutralization in mosquitoes expressing an engineered antibody.

### Permalink

<https://escholarship.org/uc/item/06h4j6xf>

### Journal

PLoS pathogens, 16(4)

### ISSN

1553-7366

### Authors

Buchman, Anna  
Gamez, Stephanie  
Li, Ming  
et al.

### Publication Date

2020-04-01

### DOI

10.1371/journal.ppat.1008545

Peer reviewed

CORRECTION

# Correction: Broad dengue neutralization in mosquitoes expressing an engineered antibody

Anna Buchman, Stephanie Gamez, Ming Li, Igor Antoshechkin, Hsing-Han Li, Hsin-Wei Wang, Chun-Hong Chen, Melissa J. Klein, Jean-Bernard Duchemin, James E. Crowe, Jr., Prasad N. Paradkar, Omar S. Akbari

In [Fig 1B](#), the columns corresponding to infection with DENV-3 and DENV-4 are duplicated and both denote the DENV-4 data. Please see the correct [Fig 1](#) here.




## OPEN ACCESS

**Citation:** Buchman A, Gamez S, Li M, Antoshechkin I, Li H-H, Wang H-W, et al. (2020) Correction: Broad dengue neutralization in mosquitoes expressing an engineered antibody. PLoS Pathog 16(4): e1008545. <https://doi.org/10.1371/journal.ppat.1008545>

**Published:** April 29, 2020

**Copyright:** © 2020 Buchman et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



**Fig 1. Effect of anti-dengue virus (DENV) single-chain variable fragment (scFv) on DENV titers of TADV-A, Wolbachia-infected (wMel), and wildtype (WT) mosquitoes.** (A) Schematic of experiment. TADV-A mosquitoes were generated via transgenesis with the anti-DENV construct, and TADV-A, wMel, and WT mosquitoes were then challenged with a blood meal infected with one of four DENV serotypes (DENV-1, isolate ET243; DENV-2, isolate ET300; DENV-3, isolate ET209; or DENV-4, isolate ET288). After the infected blood meal enters the mosquito midgut, there are two potential outcomes: in the first (applies for all tested strains), the virus replicates and disseminates past the midgut to become transmissible; in the second (applies to TADV-A mosquitoes), the anti-DENV transgene expresses scFv antibodies in the midgut that bind to the virus and neutralize it. (B) Plots depicting viral titers. To determine if the anti-DENV transgene confers resistance to all four DENV serotypes, we determined viral titers in extracted midguts, carcasses, and saliva from WT, TADV-A (homozygous [Hm] and heterozygous [Ht]), and wMel infected mosquitoes. Viral genome equivalents (GE) from mosquito midguts (at 4 days post infection [dpi]) and carcass (at 14 dpi) were determined using RT-qPCR and calculated using previously published methods. Viral titers in the saliva were determined using the median tissue culture infective dose (TCID<sub>50</sub>) on Vero cells. For each experiment, data from three replicates is pooled. Red horizontal bars represent the mean GE/viral titer. \*\* p < 0.001.

<https://doi.org/10.1371/journal.ppat.1008545.g001>

## Reference

1. Buchman A, Gamez S, Li M, Antoshechkin I, Li H-H, Wang H-W, et al. (2020) Broad dengue neutralization in mosquitoes expressing an engineered antibody. PLoS Pathog 16(1): e1008103. <https://doi.org/10.1371/journal.ppat.1008103> PMID: 31945137